



**QUEEN'S
UNIVERSITY
BELFAST**

Psychosocial interventions to improve quality of life and emotional wellbeing in newly diagnosed cancer patients

Galway, K., Black, A., Cantwell, M., Cardwell, C. R., Mills, M., & Donnelly, M. (2013). *Psychosocial interventions to improve quality of life and emotional wellbeing in newly diagnosed cancer patients*. Poster session presented at IPOS 15th World Congress of Psycho-Oncology , Rotterdam, Netherlands.
<https://doi.org/10.1002/14651858.CD007064.pub2>

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
Copyright the authors 2013.

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.



Psychosocial interventions to improve quality of life and emotional wellbeing for recently diagnosed cancer patients

Galway K⁺, Black A^s, Cantwell M⁺, Cardwell CR⁺, Mills M[#], Donnelly M⁺



⁺Queens University Belfast, NI ^s National Cancer Institute, Bethesda, USA [#] Northern Health and Social Care Trust, NI

INTRODUCTION

Cancer occurs in one in four of the population with over a quarter of a million people in the UK diagnosed each year (excluding non-melanoma skin cancer).

A diagnosis of cancer can be emotionally challenging. **UK government policy recommends** that all individuals who are diagnosed with cancer should be assessed for emotional problems and given **access to appropriate psychological support services**. However, evidence to guide the nature and content of such services and their mode of delivery is contradictory.

This review examines the effectiveness of individual psychosocial interventions in the first 12 months after diagnosis. **Psychosocial interventions are defined** as involving a 'trained helper' providing therapeutic dialogue, sometimes referred to as **talking therapy**, with an individual diagnosed with cancer with the aim of improving **quality of life and emotional wellbeing**.

SUMMARY OF FINDINGS

Psychosocial interventions to improve quality of life and emotional wellbeing compared with usual care for newly diagnosed cancer patients

Patient or population: newly diagnosed patients with cancer
Settings: hospital and community based settings
Intervention: psychosocial interventions
Comparison: usual care

Outcomes	Standardised Mean Differences (95% CI)	No of Participants (studies)
Measures of Quality of Life [Psychometric measures of quality of life] [6 months]	0.11 [-0.00, 0.22]	1249 (9)
General Psychological Distress [Psychometric measures of general psychological distress] [6 months]	0.08 [-0.05, 0.21]	1502 (9)
Mood Measures [Psychometric measures of mood] [6 months]	-0.81 [-1.44, -0.18]	683 (8)

The results are **inconclusive** with much heterogeneity across studies

No improvement in general quality of life was found, but **small improvements in 'illness related' quality of life** were observed

No improvements in anxiety or depression were found, but **small improvements in mood were detected**

Nurse-led interventions, along with **psycho-educational** interventions using **telephone and face-to-face delivery** appear to show some promise

METHODS – RCTs - PICO

PARTICIPANTS

Recently diagnosed with cancer

INTERVENTIONS

Individual, interpersonal dialogue with a trained helper


COMPARISON

Usual care


OUTCOMES

Primary outcome = Quality of life (QoL)

Secondary outcomes = General psychological distress (including depression and anxiety) and mood



RISK OF BIAS GRAPH



Inconsistent reporting gave rise to unclear risk of bias (above) across many studies

FUTURE RECOMMENDATIONS

- **Improve the quality of reporting** in psychosocial RCT publications, to facilitate future syntheses of results and reduce risk of bias
- **More work is needed to match types of patients to types of psychosocial interventions** e.g. using a needs led approach
- **Use appropriate psychometrics for the trial population** and avoid measures of clinical diagnoses in non-clinical populations
- **Evaluate which type of 'trained helper' is the most appropriate** professional to deliver psychosocial interventions
- **Conduct economic appraisals** of the cost-effectiveness of interventions
- **Synthesise patient reported preferences** reported in qualitative publications

Reference: Galway K, Black A, Cantwell M, Cardwell CR, Mills M, Donnelly M. Psychosocial interventions to improve quality of life and emotional wellbeing for recently diagnosed cancer patients. Cochrane Database of Systematic Reviews 2012, Issue 11. Art. No.: CD007064. DOI: 10.1002/14651858.CD007064.pub2.